

Ground State Phase Diagram of Frustrated $S = 1$ XXZ chains : Chiral Ordered Phases

Toshiya HIKIHARA*, Makoto KABURAGI¹, Hikaru KAWAMURA² and Takashi TONEGAWA³

Graduate School of Science and Technology, Kobe University, Rokkodai, Kobe 657-8501

¹*Faculty of Cross-Cultural Studies, Kobe University, Tsurukabuto, Nada, Kobe 657-8501*

²*Faculty of Engineering and Design, Kyoto Institute of Technology, Sakyo-ku, Kyoto 606-8585*

³*Department of Physics, Faculty of Science, Kobe University, Rokkodai, Kobe 657-8501*

(Received September 16, 1999)

The ground-state phase diagram of frustrated $S = 1$ XXZ spin chains with the competing nearest- and next-nearest-neighbor antiferromagnetic couplings is studied using the infinite-system density-matrix renormalization-group method. We find six different phases, namely, the Haldane, gapped chiral, gapless chiral, double Haldane, Néel, and double Néel (uudd) phases. The gapped and gapless chiral phases are characterized by the spontaneous breaking of parity, in which the long-range order parameter is a chirality, $\kappa_l = S_l^x S_{l+1}^y - S_l^y S_{l+1}^x$, whereas the spin correlation decays either exponentially or algebraically. These chiral ordered phases appear in a broad region in the phase diagram for $\Delta < 0.95$, where Δ is an exchange-anisotropy parameter. The critical properties of phase transitions are also studied.

KEYWORDS: frustration, spin-1 chain, ground-state phase diagram, density-matrix renormalization-group method, chiral ordering

§1. Introduction

Frustrated antiferromagnetic quantum spin systems have attracted considerable attention over the decades since they exhibit a wealth of fascinating phenomena in their ground states and low-lying excitations. In general, frustration suppresses antiferromagnetic correlations and the tendency towards the Néel order. Classical systems, for example, often show a helical ordered state in their ground states in the presence of strong frustration. In quantum systems, the interplay of frustration and quantum fluctuations plays an important role which causes exotic phenomena, *e.g.*, a spin-liquid state and a novel type of spontaneous symmetry breaking.

In this paper, we study a one-dimensional anisotropic spin system with the antiferromagnetic nearest-neighbor coupling J_1 and the frustrating next-nearest-neighbor coupling J_2 . The model is

* E-mail address: hikihara@phys03.phys.sci.kobe-u.ac.jp

described by the XXZ Hamiltonian

$$\mathcal{H} = \sum_{\rho=1}^2 \left\{ J_{\rho} \sum_l \left(S_l^x S_{l+\rho}^x + S_l^y S_{l+\rho}^y + \Delta S_l^z S_{l+\rho}^z \right) \right\}, \quad (1.1)$$

where $\mathbf{S}_l$ is a spin- S spin operator at site l and $\Delta \geq 0$ represents an exchange anisotropy. Hereafter, we put $j \equiv J_2/J_1$ ($j \geq 0$).

In the classical limit, $S \rightarrow \infty$, the system exhibits a magnetic long-range order (LRO) characterized by a wavenumber q . The order parameter is defined by

$$\mathbf{m}(q) = \frac{1}{L} \sum_l \mathbf{S}_l e^{iq l}, \quad (1.2)$$

where L is the total number of spins. While the magnetic LRO is of the Néel-type ($q = \pi$) when j is smaller than a critical value, $j \leq 1/4$, it becomes of helical-type for $j > 1/4$ with $q = \cos^{-1}(-1/4j)$. It should be noticed that both the time-reversal and parity symmetries are broken in this helical ordered state. When the system has an XY -like anisotropy ($\Delta < 1$), the helical ordered state possesses a two-fold discrete degeneracy according as the helix is either right- or left-handed, in addition to a continuous degeneracy associated with the original $U(1)$ symmetry of the XY spin. This discrete degeneracy is characterized by mutually opposite signs of the total chirality defined by,¹⁾

$$O_{\kappa} = \frac{1}{L} \sum_l \kappa_l, \quad (1.3)$$

$$\kappa_l = S_l^x S_{l+1}^y - S_l^y S_{l+1}^x = [\mathbf{S}_l \times \mathbf{S}_{l+1}]_z.$$

Note that this chirality is distinct from the scalar chirality of the Heisenberg spin often discussed in the literature²⁾ defined by $\chi_l = \mathbf{S}_{l-1} \cdot \mathbf{S}_l \times \mathbf{S}_{l+1}$: The chirality O_{κ} changes its sign under the parity operation but is invariant under the time-reversal operation while the scalar chirality changes its sign under both operations. Since the frustrated classical chain (1.1) always has a planar spin order, the scalar chirality χ_l vanishes trivially.

In the $S = 1/2$ case, the ground-state phase diagram of the system has been extensively studied either numerically^{3,4,5,6)} or analytically.^{7,8)} These studies have revealed that the system undergoes a phase transition from the spin-liquid phase to the dimer phase at $j = j_c(\Delta)$ with increasing j . The critical value $j_c(\Delta)$ has been estimated to be $j_c \sim 0.241$ for the Heisenberg case ($\Delta = 1$).^{4,5,6)} The spin-liquid phase at $j \leq j_c(\Delta)$ is characterized by gapless excitations and an algebraic decay of spin correlations, while the dimer phase at $j > j_c(\Delta)$ is characterized by a finite energy gap above the doubly degenerate ground states and by the spontaneous breaking of both the parity and translational symmetries. The dimer phase is divided into two regions by the Lifshitz line $j_L(\Delta)$ according to the nature of the short-range spin correlations, where $j_L = 0.5$ in the Heisenberg case.⁹⁾ Whereas the correlations are of the Néel type and the structure factor $S(q)$ has a maximum

at $q = \pi$ for $j \leq j_L(\Delta)$, the correlations become incommensurate and the maximum of $S(q)$ occurs at $q = Q < \pi$ for $j > j_L(\Delta)$. This incommensurate character is regarded as the vestige of the helical order in the classical case. It has also been shown both in the XY and Heisenberg cases that there is no chiral LRO for arbitrary j , i.e., the chirality O_κ vanishes in the thermodynamic limit.¹⁰⁾

As is well known, on the other hand, the low-energy properties of the $S = 1$ spin chain without frustration are essentially different from those of the $S = 1/2$ chain. Haldane conjectured that integer spin chains are in a phase with a finite energy gap in the excitation spectrum and with an exponential decay of the two-spin correlation functions, while half-odd integer spin chains are in the spin-liquid phase.^{11,12)} This gapped phase for integer spin chains is now called the Haldane phase, and is characterized by the string order parameter¹³⁾

$$O_{\text{str}} = \frac{1}{L} \sum_l \exp \left(\sum_{j=1}^{l-1} i\pi S_j^z \right) S_l^z. \quad (1.4)$$

Haldane's conjecture has been verified either numerically or experimentally, and is now well-established. It is therefore natural to expect that the frustrated $S = 1$ spin chain might also exhibit low-energy properties which are substantially different from those of both the $S = 1/2$ and the $S = \infty$ classical chains.

The aim of this paper is to determine the ground-state phase diagram of the model (1.1) with $S = 1$ spins in the j - Δ plane. The ground-state properties of the model have been studied in some limiting cases. In the Heisenberg case ($\Delta = 1$), Kolezhuk *et al.* have shown that, at $j = 0.7444(6)$, the system undergoes a first-order phase transition from the Haldane phase to the 'double Haldane' (DH) phase, in which the next-nearest-neighbor coupling J_2 is dominant and the system can be regarded as two Haldane subchains coupled with a weak inter-subchain coupling J_1 .^{14,15)} The string order parameter O_{str} vanishes discontinuously at the transition. Meanwhile, in the case of no frustration $j = 0$, it has been shown that a phase transition from the Haldane phase to the Néel phase occurs at $\Delta = \Delta_c \sim 1.18$.^{16,17,18,19)} The transition in this limit has been verified to belong to the universality class of the two-dimensional (2D) Ising model. It is also known that, in the Ising limit, $\Delta \rightarrow \infty$, there is a phase transition at $j = 0.5$ between the Néel phase and the 'double Néel' (DN) phase, the latter being characterized by the magnetic LRO of 'up-up-down-down' ('uudd') type. Systematic studies of the model in the entire j - Δ plane, however, has been lacking so far.

In the previous work,¹⁰⁾ we have shown that in the pure XY case ($\Delta = 0$) there occurs a phase transition from the Haldane phase to the 'chiral ordered' phase in which only the chirality exhibits a LRO ($O_\kappa > 0$) with no helical LRO ($\mathbf{m}(q) = 0$ for $\forall q$). Here we note that, in the pure XY and Heisenberg cases, the absence of magnetic (spin) LRO has rigorously been proven for any j and for general $S < \infty$.²⁰⁾ We have also found that there are two types of the chiral phase, the 'gapped' and 'gapless' chiral phases, in which the spin correlation decays exponentially and algebraically,

respectively. These chiral phases spontaneously break only the parity symmetry, with preserving both the time-reversal and translational symmetries. The phase defined by this types of symmetry breaking is a novel one in quantum systems, while it has been studied quite intensively in various frustrated classical systems.^{21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31)} It is thus interesting to further investigate the properties of the chiral phases in frustrated quantum systems.

In order to determine the phase diagram, we numerically calculate the correlation functions associated with the order parameters, $\mathbf{m}(q)$, O_κ , and O_{str} , for open chains, using the infinite-system density-matrix renormalization-group (DMRG) method. Examining the behavior of these correlation functions at long distance, we find the following six phases, the Haldane, gapped and gapless chiral, DH, Néel, and DN phases, in the region where $j \geq 0$ and $\Delta \geq 0$. The obtained phase diagram has revealed that the gapped and gapless chiral phases exist in a broad region of the diagram where $0 \leq \Delta < 0.95$.

The plan of this paper is as follows. In § 2, we introduce the spin, chiral, and string correlation functions associated with each order parameter. We also explain our numerical method in some detail. Results of our numerical calculations are presented in § 3. The ground-state phase diagram in the j - Δ plane is constructed and the critical properties of the phase transitions between different phases are analyzed. Finally, we summarize our results in § 4.

§2. Correlation Functions and Numerical Method

In order to determine the phase diagram, we calculate the spin, chiral, and string correlation functions defined by,

$$C_s^\alpha(r) = \langle S_{l_0-r/2}^\alpha S_{l_0+r/2}^\alpha \rangle \quad (\alpha = x, z), \quad (2.1)$$

$$C_\kappa(r) = \langle \kappa_{l_0-r/2} \kappa_{l_0+r/2} \rangle, \quad (2.2)$$

$$C_{\text{str}}(r) = \langle S_{l_0-r/2}^z \exp \left(i\pi \sum_{j=l_0-r/2}^{l_0+r/2-1} S_j^z \right) S_{l_0+r/2}^z \rangle, \quad (2.3)$$

which are associated with the helical, chiral, and string order parameters, respectively. Here, l_0 represents the center position of the open chain, i.e., $l_0 = L/2$ for even r and $l_0 = (L+1)/2$ for odd r , and $\langle \cdots \rangle$ represents the expectation value in the lowest energy state in the subspace of $S_{\text{total}}^z = 0$ with even parity. We have checked by exact-diagonalization calculation that the ground state of the system indeed belongs to this subspace. We employ the infinite-system DMRG algorithm introduced by White.^{32, 33)} The algorithm is accelerated by using the initial Lanczos vector derived from the recursion relation proposed by Nishino and Okunishi.^{34, 35)}

Here we note the following two points concerning the properties of the infinite-system DMRG. The first is the dependence of the data on the number of kept states, m . Since no analytic form to extrapolate the data to $m \rightarrow \infty$ has been established so far, in order to obtain precise results we have no choice but to increase m until the data converge. Generally speaking, the convergence

is slow when the energy gap is small, while it is fast when the gap is large. In fact, we have kept up to $m = 400$ states in the vicinity of phase boundaries and in the gapless chiral phase, whereas $m = 220$ turns out to be large enough in the gapped phases.

Besides, we need to check the L -convergence of the correlations. In the infinite-system DMRG, two spins are added in the center of the chain at each step of the algorithm. We continue the procedure until the correlation functions calculated around the center become free from the effects of open boundaries. In general, more steps are needed if the system has longer correlations. We have performed calculations with up to 3500 steps, which corresponds to the system of $L = 7000$, until the L -convergence of the data of the correlation functions was attained.

§3. Numerical Results

Examining the r -dependence of the correlation functions given by eqs.(2.1)-(2.3), we find six different phases in the phase diagram, namely, the Haldane, gapped chiral, gapless chiral, DH, Néel, and DN phases. Figure 1 shows the obtained phase diagram in the j - Δ plane. The long-distance behavior of the correlation functions in each phase is summarized in Table I. In the following subsections, we discuss our numerical results at each transition line in more detail.

3.1 Transition to the chiral phases

Let us first consider the phase transition between the Haldane and chiral ordered phases. In our previous work,¹⁰⁾ we studied the phase transition in the pure XY case ($\Delta = 0$), and found that there exist two different types of chiral ordered phase. In the gapped chiral phase, the chiral and string LRO's coexist and the spin correlations decay exponentially with a finite energy gap, while in the gapless chiral phase, the chiral LRO exists and both the spin and string correlations decay algebraically with gapless excitations. We also showed that the gapped chiral phase existed in a very narrow region of j between the Haldane and the gapless chiral phases. As j increases, two transitions, the Haldane-gapped chiral transition at $j = j_{c1}(\Delta = 0) = 0.473 \pm 0.001$ and the gapped chiral-gapless chiral transition at $j = j_{c2}(\Delta = 0) = 0.490^{+0.010}_{-0.005}$, occur successively.

In the present calculation, we find that the gapped chiral phase continues to exist for $\Delta > 0$. As an example, we show in Fig. 2 the r -dependence of the spin, chiral, and string correlations for $\Delta = 0.4$ and several typical values of j on log-log plots. One can clearly see from the figure, both the chiral and string LRO's coexist at $j = 0.590$, which indicates that the point $j = 0.590$ belongs to the gapped chiral phase. Performing the calculations with increasing j at intervals of 0.002, we estimate the transition points for $\Delta = 0.4$ to be $j_{c1} = 0.582 \pm 0.002$ and $j_{c2} = 0.600 \pm 0.003$.

Since j_{c1} lies so close to j_{c2} , one may wonder if the appearance of the gapped chiral phase might be just an artifact of the numerical calculation. We, however, believe that we can refute such suspicion. In our infinite-system DMRG, the only source of uncertainty comes from the truncation error due to a finite m . As can be seen in Fig. 2, both the chiral correlations in the Haldane phase

($j \lesssim 0.580$) and the string correlations in the gapless chiral phase ($j \gtrsim 0.600$) tend to be enhanced with increasing m . This means that the estimated transition point j_{c1} (j_{c2}) monotonously decreases (increases) as m is increased, i.e., the DMRG calculation tends to underestimate the range of j for the gapped chiral phase. We can therefore safely conclude that the two transition points locate very near, but are distinctly different from each other, and the gapped chiral phase does exist. Analyzing the data for several Δ , we find the gapped chiral phase in the range $0 \leq \Delta \leq 0.8$. The estimated values of $j_{c1}(\Delta)$ and $j_{c2}(\Delta)$ are summarized in Table II. At $\Delta = 0.9$, no region belonging to the gapped chiral phase is observed within the calculation with varying j at intervals of 0.002. We, however, cannot precisely determine the point beyond which the gapped chiral phase vanishes, since we cannot exclude the possibility that the region of the gapped chiral phase is just too narrow to detect even at $\Delta = 0.9$.

In order to investigate the critical properties of the Haldane-gapped chiral phase transition at $j = j_{c1}$, we analyze the chiral correlation length ξ_κ in the Haldane phase. The chiral correlation length is estimated by fitting the chiral correlation functions in the range $30 \lesssim r \leq 100$ to the standard form of an exponential decay,

$$C_\kappa(r) = A \exp(-r/\xi_\kappa), \quad (3.1)$$

where A is a constant. The j -dependence of the estimated chiral correlation length ξ_κ in the pure XY case ($\Delta = 0$) is shown in Fig. 3 on a log-log plot. It shows a linear behavior near the critical point suggesting the standard power-law divergence, $\xi_\kappa \sim |j - j_{c1}|^{-\nu_\kappa}$. By least-square fitting using the three data points near the critical point, we estimate $\nu_\kappa = 0.9 \pm 0.1$. The error is rather large since the exponent is sensitive to the change of the critical point j_{c1} . The estimated value of ν_κ is slightly smaller than, but consistent with, that of the 2D Ising model, $\nu_\kappa = 1$.

At the gapped chiral-gapless chiral transition at $j = j_{c2}$, we extract the exponents of algebraic decay of the spin correlation function $C_s^x(r)$ in the gapless chiral phase by fitting $C_s^x(r)$ in the range $10 \lesssim r \leq 100$ to the power-law form,

$$C_s^x(r) = B r^{-\eta_x}, \quad (3.2)$$

where B is a constant. Figure 4 shows the j -dependence of the estimated exponent η_x in the pure XY case ($\Delta = 0$). As can be seen in the figure, the exponent η_x increases from $\eta_x \sim 0.15$ at $j \gtrsim 0.8$ to $\eta_x \sim 0.29$ at $j = 0.490$ as j approaches the critical value, $j = j_{c2} = 0.490_{-0.005}^{+0.010}$. We estimate η_x at the critical point to be $\eta_x(j = j_{c2}) = 0.28 \pm 0.03$. The large error is due to the rather large error in the estimated value of j_{c2} . Although the estimated value of η_x at $j = j_{c2}$ is not inconsistent with the value for the Kosterlitz-Thouless transition, $\eta_x = 1/4$, more detailed study is needed to identify the universality class of the transition between the gapped and gapless chiral phases.

Next, we consider the transition between the chiral and DH phases at $j = j_{c3}$. As mentioned in § 1, the system in the DH phase can be regarded as two Haldane subchains with the exchange

coupling J_2 coupled with the weak inter-subchain coupling J_1 . Thus it is natural to expect the string correlations on each subchain to be the order parameter of this phase. This generalized string correlation, however, does not exhibit a LRO in this phase, but decays exponentially.^{14,15} Consequently, we have no order parameter at hand to characterize the DH phase. We accordingly determine the phase boundary between the gapless chiral and DH phases as the line where the chiral LRO vanishes. The data of the chiral correlation for $j = 1.0$ and several typical values of Δ are shown in Fig. 5. It is clearly shown in the figure that the system exhibits a chiral LRO for $\Delta \lesssim 0.68$ while $C_\kappa(r)$ decays exponentially for $\Delta \geq 0.70$. Hence, we estimate the transition point, $j_{c3}(\Delta = 0.69 \pm 0.01) = 1.0$. As can be seen in Fig.1, the value of $j_{c3}(\Delta)$ estimated in this way becomes larger as Δ becomes smaller. Here we note that there is a possibility that the gapped chiral phase also exists in the vicinity of the line, i.e., the line where the chiral LRO vanishes might be different from the line where the energy gap vanishes. It is, however, very difficult to distinguish these lines each other, if any, since we have no means to estimate the energy gap accurately enough to do this. For that purpose, it may be necessary to introduce another order parameter which characterizes the DH phase.

Since the z -component of the exchange couplings in the Hamiltonian, $S_i^z S_{i+\rho}^z$, reduces the energy barrier between the two discretely degenerate states according to the right- and left-handed chirality, we can expect naively that the tendency towards chiral ordering is suppressed by the introduction of the z -component. This feature can clearly be seen in Fig.1, i.e., the chiral phases appear in the narrower range of j as Δ becomes larger. While the chiral LRO is found for $0 \leq \Delta \leq 0.9$ in some range of j , we observe no chiral LRO for $\Delta = 0.95$ and 1.0 . From this observation, we conclude that, between $0.9 < \Delta < 0.95$, there is a multi-critical point among the Haldane, gapless chiral, and DH phases.

3.2 Haldane-DH transition

Here we consider the transition between the Haldane and DH phases at $j = j_{c4}$. This transition has been studied by Kolezhuk *et al.* in the Heisenberg case ($\Delta = 1$).^{14,15} From the numerical calculation of the string correlation function $C_{\text{str}}(r)$ by the DMRG method, they estimated the transition point, $j = j_{c4} = 0.7444(6)$, where the string LRO vanished. They also concluded that the transition was first-order from the observation of a discontinuous jump of the string order parameter from a finite value to zero. Our results are consistent with theirs; we have also observed the jump in the string order parameter at the transition, although the estimated transition point $j = j_{c4} = 0.747 \pm 0.001$ is slightly different from theirs. The reason of this small difference may be that the number of kept states, m , of our calculation are larger than that of their calculation. The first-order nature of the phase transition persists even away from the Heisenberg line ($\Delta = 1$). In Fig. 6, we show the j -dependence of the string correlation at $r \rightarrow \infty$ for $\Delta = 0.95$ and 1.20 . Clearly, the string order parameter disappears discontinuously at the transition point. The

estimated transition line runs from the multi-critical point to the Néel-DN phase transition point, $j = 0.5$ in the Ising limit $\Delta \rightarrow \infty$, which will be explained in the next subsection.

3.3 Transition to the magnetic ordered phases

In this subsection, we discuss the results for the Haldane-Néel transition at $j = j_{c5}$ and the region between the DH and DN phases. Some information is available at some special points in the phase diagram. In the Ising limit, $\Delta \rightarrow \infty$, a phase transition between the Néel and the DN phases occurs at $j = 0.5$. The system exhibits a magnetic LRO in both phases while the nature of two-spin correlations changes from the Néel type ($j \leq 0.5$) to the ‘uudd’ type ($j > 0.5$). In the case of $j = 0$, it has also been shown analytically^{16,17)} and numerically^{18,19)} that there is a phase transition between the Haldane and Néel phases at $\Delta = \Delta_c \sim 1.18$. This transition is continuous and belongs to the universality class of the 2D Ising model. In the limit of $j \rightarrow \infty$, since the system can be regarded as two decoupled chains with the exchange coupling J_2 , there must be a phase transition of the same universality class between the DH and DN phases at $\Delta = \Delta_c$.

We estimate numerically the phase boundary between the Haldane and Néel phases by examining the existence of the magnetic LRO of the Néel type. As an example, we show in Fig. 7 our numerical results of $C_s^z(r)$ for $\Delta = 1.5$ and several typical values of j , which suggests that the transition occurs at $j = j_{c5}(\Delta = 1.5) \sim 0.1845$. As can be seen in Fig. 1, the phase boundary determined in this way smoothly connects the transition point in the limiting case, $\Delta = \Delta_c$ for $j = 0$, to the Néel-DN transition point, $j = 0.5$ for $\Delta \rightarrow \infty$. Since the Haldane-Néel transition at $j = 0$ belongs to the universality class of the 2D Ising type, we expect that the transition in the intermediate region also exhibits the same critical properties. We analyze the critical behavior of the phase transition in the intermediate region at $\Delta = 1.5$. From the analysis of the order parameter and the correlation length associated with the z -component of the spin shown in Figs.8(a) and 8(b), we obtain the critical exponents, $\beta = 0.122 \pm 0.010$ and $\nu = 0.96 \pm 0.08$. We also estimate the critical-point decay exponent $\eta = 0.26 \pm 0.02$ from the data of $C_s^z(r)$ shown in Fig. 7. These values of critical exponents are in good agreement with the values of the 2D Ising model, $\beta = 1/8$, $\eta = 1/4$ and $\nu = 1$.

Next we consider the region between the DH and DN phases. As an example, we show in Fig. 9 the numerical data of $C_s^z(r)$ for $j = 1.0$ and several typical values of Δ on a semi-log plot. In the DH phase ($\Delta \leq 1.6$), the spin correlation turns out to be incommensurate (note that $C_s^z(r)$ in this phase has been divided by $\cos(Qr)$ in the figure) and decays exponentially, while it exhibits a magnetic LRO of the ‘uudd’ type in the DN phase ($\Delta \geq 2.0$). In the DH phase, the correlation length associated with the incommensurate short-range order becomes longer as Δ increases. For $\Delta = 1.7, 1.8$, and 1.9 , on the other hand, our infinite-system DMRG calculation does not converge, i.e., the bond energy at the center of the chain fluctuates irregularly even after thousands of steps of the algorithm. This means that in the region it is impossible to obtain the bulk properties of the system within the framework of the infinite-system DMRG method. The origin of this difficulty

can be attributed to an approximation employed in the algorithm. As noted in § 2, the size of the system treated in the infinite-system DMRG increases by two at every step. Accordingly, the set of m -states kept in a step is, strictly speaking, not the optimal one for the block states in the next step. In the absence of magnetic LRO or in the presence of commensurate LRO with the periodicity of two or four lattice sites ($q = \pi$ or $\pi/2$), this error of the algorithm becomes less and less significant as the system becomes larger, while in the presence of incommensurate (or highercommensurate) magnetic LRO with a wavenumber $q = Q \neq \pi, \pi/2$, the error remains even in the thermodynamic limit and causes a crucial problem in the calculation.³⁶⁾ We therefore consider that the breakdown of our calculation in the range $1.65 \lesssim \Delta \lesssim 1.95$ suggests that the system has an incommensurate magnetic LRO in this region. From this conjecture, we believe that the point where the magnetic LRO of the ‘uudd’ type starts to appear, $j = 1.0$ and $\Delta \sim 1.95$ (see Fig. 9), is a boundary point of the DN phase. Connecting the points estimated in this way, we determine the boundary line of the DN phase (the dashed line in Fig. 1). In contrast, we cannot precisely determine the boundary line of the DH phase nor directly verify the existence of the phase with an incommensurate magnetic LRO.

§4. Summary

In this paper we have studied the ground-state properties of spin-1 XXZ chains with the antiferromagnetic nearest-neighbor and the frustrating next-nearest-neighbor couplings. We have performed the infinite-system DMRG calculation to numerically compute the spin, chiral, and string correlation functions. Analyzing the r -dependence of these correlations, we have determined the ground-state phase diagram which turns out to include the six phases, the Haldane, gapped and gapless chiral, DH, Néel, and DN phases (Fig.1). The chiral phases appear in a quite broad region in the phase diagram for $\Delta < 0.95$, while the region becomes narrower as Δ increases. We have also confirmed that the gapped chiral phase does exist in a very narrow but finite region between the Haldane and gapless chiral phases. The multi-critical point among the Haldane, gapless chiral, and DH phases exists between $0.9 < \Delta < 0.95$. Furthermore, we have investigated the critical properties of the phase transitions. For the Haldane-gapped chiral transition, the chiral correlation-length exponent is estimated to be $\nu_\kappa = 0.9 \pm 0.1$, which is not inconsistent with the 2D Ising value, $\nu_\kappa = 1$.

The quantum chiral phases which we have discovered in this and previous work are novel ones in quantum spin systems. Accordingly, many interesting open problems about these phases still remain. As mentioned in § 1, in the $S = 1$ case only the chirality exhibits a LRO without the helical LRO in the chiral phases while both LRO coexist in the helical ordered phase which appears in the classical case, $S \rightarrow \infty$. This means that the quantum fluctuation in the $S = 1$ chain recovers only the time-reversal symmetry with the parity symmetry kept broken. In the $S = 1/2$ case, strong quantum fluctuation eventually recovers both symmetries. In order to gain more insight into the

effect of quantum fluctuation on each symmetry, it is necessary to investigate how the phase diagram and the nature of the chiral phases depend on the spin quantum number S , particularly paying attention to the possible difference between integer and half-odd integer S . We are now performing calculations in the $S = 3/2$ case. A preliminary calculation suggests that the gapless chiral phase appears in this case as well.

Acknowledgments

We thank K. Nomura and T. Momoi for useful discussions. Numerical calculations were carried out in part at the Yukawa Institute Computer Facility, Kyoto University. T. H. was supported by a Grant-in-Aid for Encouragement of Young Scientists from Ministry of Education, Science and Culture of Japan.

-
- [1] H. Kawamura: Phys. Rev. B **38** (1988) 4916.
 - [2] H. Frahm and C. Rödenbeck: J. Phys. A **30** (1997) 4467.
 - [3] I. Harada and T. Tonegawa: *Recent Advances in Magnetism of Transition Metal Compounds*, ed. A. Kotani and N. Suzuki (World Scientific, Singapore, 1993) p. 348.
 - [4] K. Nomura and K. Okamoto: J. Phys. A **27** (1994) 5773.
 - [5] K. Okamoto and K. Nomura: Phys. Lett. A **169** (1992) 433.
 - [6] K. Nomura and K. Okamoto: J. Phys. Soc. Jpn. **62** (1993) 1123.
 - [7] C.K. Majumder and D.K. Ghosh: J. Math. Phys. **10** (1969) 1399.
 - [8] F.D.M. Haldane: Phys. Rev. B **25** (1982) 4925.
 - [9] T. Tonegawa and I. Harada: J. Phys. (Paris) Suppl. **49** (1988) C8-1411.
 - [10] M. Kaburagi, H. Kawamura and T. Hikihara: cond-mat/9902002, to appear in J. Phys. Soc. Jpn.
 - [11] F. D. M. Haldane: Phys. Lett. **93A** (1983) 464.
 - [12] F. D. M. Haldane: Phys. Rev. Lett. **50** (1983) 1153.
 - [13] M. den Nijs and K. R. Rommelse: Phys. Rev. B **40** (1989) 4709.
 - [14] A. Kolezhuk, R. Roth and U. Schollwöck: Phys. Rev. Lett. **77** (1996) 5142.
 - [15] A. Kolezhuk, R. Roth and U. Schollwöck: Phys. Rev. B **55** (1997) 8928.
 - [16] G. Gómez-Santos: Phys. Rev. Lett. **63** (1989) 790.
 - [17] G. Gómez-Santos: J. Appl. Phys. **67** (1990) 5610.
 - [18] J. C. Bonner: J. Appl. Phys. **61** (1987) 3941.
 - [19] K. Nomura: Phys. Rev. B **40** (1989) 9142.
 - [20] T. Momoi: J. Stat. Phys. **85** (1996) 193.
 - [21] H. Kawamura: J. Phys. Condens. Matter **10** (1998) 4707.
 - [22] S. Miyashita and H. Shiba: J. Phys. Soc. Jpn. **53** (1984) 1145.
 - [23] D. H. Lee, J.D. Jannopoulos, J. W. Negele and D. P. Landau: Phys. Rev. Lett. **52** (1984) 433.
 - [24] L. Capriotti, R. Vaia, A. Cuccoli and V. Tognetti: Phys. Rev. B **58** (1998) 273.
 - [25] S. Teitel and C. Jayaprakash: Phys. Rev. B **27** (1983) 598.
 - [26] G. Ramirez-Santiago and J. V. José: Phys. Rev. Lett. **68** (1992) 1224.
 - [27] P. Olsson: Phys. Rev. Lett. **75** (1995) 2758.
 - [28] P. Olsson: Phys. Rev. B **55** (1997) 3585.
 - [29] H. Kawamura and M. Tanemura: J. Phys. Soc. Jpn. **60** (1991) 1092.

- [30] H. Kawamura: Phys. Rev. B **51** (1995) 12398.
- [31] H. Kawamura: Phys. Rev. Lett. **82** (1999) 964.
- [32] S. R. White: Phys. Rev. Lett. **69** (1992) 2863.
- [33] S. R. White: Phys. Rev. B **48** (1993) 10345.
- [34] T. Nishino and K. Okunishi: J. Phys. Soc. Jpn. **64** (1995) 4084.
- [35] T. Hieida, K. Okunishi and Y. Akutsu: Phys. Lett. A **233** (1997) 464.
- [36] See, for example, *Density-Matrix Renormalization*, ed. I. Peschel, X. Wang, M. Kaulke and K. Hallberg (Springer Verlag, Berlin, 1999).

Table I. The long-distance behavior of the spin, chiral, and string correlations in each phase. "expo." and "power" means exponential decay and power-law decay, respectively.

	Haldane	gapped chiral	gapless chiral	DH	Néel	DN
C_s^z	expo.	expo.	power	expo.	LRO	LRO
C_s^x	expo.	expo.	power	expo.	expo.	expo.
C_κ	expo.	LRO	LRO	expo.	expo.	expo.
C_{str}	LRO	LRO	power	expo.	LRO	LRO

Table II. The estimated values of $j_{c1}(\Delta)$ and $j_{c2}(\Delta)$ for various values of Δ .

Δ	0	0.2	0.4	0.6	0.8
j_{c1}	0.473 ± 0.001	0.527 ± 0.002	0.582 ± 0.002	0.642 ± 0.002	0.715 ± 0.003
j_{c2}	$0.490^{+0.010}_{-0.005}$	0.540 ± 0.003	0.600 ± 0.003	0.660 ± 0.003	0.720 ± 0.002

Fig. 1. The ground-state phase diagram. The circles, squares, and diamonds represent the points where the string LRO, chiral LRO, and magnetic LRO vanishes, respectively.

Fig. 2. The r -dependence of the correlation functions for $\Delta = 0.4$: (a) spin correlation $C_s^x(r)$ divided by the oscillating factor $\cos(Qr)$; (b) chiral correlation $C_\kappa(r)$; (c) string correlation $-C_{\text{str}}(r)$. The number of kept states is $m = 260$ (for $j = 0.570$), $m = 300$ (for $j = 0.580, 0.590, 0.610$) and $m = 400$ (for $j = 0.600$). To illustrate the m -dependence, we also indicate the data with smaller m by crosses for several cases: in figure (b), $m = 180, 220$ for $j = 0.570$ and $m = 220, 260$ for $j = 0.580$; in figure (c), $m = 300, 350$ for $j = 0.600$ and $m = 260, 300$ for $j = 0.610$. In the other cases, the numerical errors of the data are smaller than the symbols. Arrows in the figures represent the extrapolated $r = \infty$ values.

Fig. 3. The log-log plot of the j -dependence of the chiral correlation length for $\Delta = 0$ (XY case) in the critical region of the Haldane phase ($j < j_{c1}$). We plot the data setting $j_{c1} = 0.4729$.

Fig. 4. The j -dependence of the exponent of algebraic decay of $C_s^x(r)$ for $\Delta = 0$ in the gapless chiral phase ($j > j_{c2} \sim 0.490$). The vertical dashed line represents the phase boundary j_{c2} .

Fig. 5. The r -dependence of the chiral correlation function for $j = 1.0$ on a log-log plot. The number of kept states is $m = 300$ (for $\Delta = 0.60, 0.65, 0.72$), $m = 350$ (for $\Delta = 0.68$) and $m = 400$ (for $\Delta = 0.70$). For $\Delta = 0.68$, where the m -dependence of the data is rather large, we indicate the data with smaller m by crosses ($m = 260, 300$). For the other Δ , the numerical errors of the data are smaller than the symbols. Arrows in the figures represent the extrapolated $r = \infty$ values.

Fig. 6. The strength of the string correlation function at $r \rightarrow \infty$ as a function of j for $\Delta = 0.95$ (solid circles) and 1.20 (open circles). The vertical dashed lines represent the transition points j_{c4} for each Δ .

Fig. 7. The r -dependence of the spin correlation function $C_s^z(r)$ for $\Delta = 1.5$ on a log-log plot. The number of kept states is $m = 220$ for all j . The numerical errors of the data are smaller than the symbols. The solid line represents the power-law decay of the correlation at $j = 0.1845$.

Fig. 8. (a) The strength of the spin correlation function $C_s^z(r)$ at $r \rightarrow \infty$ in the Néel phase ($j < j_{c5}$) as a function of j for $\Delta = 1.5$. We plot the data setting $j_{c5} = 0.1845$. We extract the exponent β by fitting the data to the form, $C_s^z(r \rightarrow \infty) \sim |j - j_{c5}|^{2\beta}$. (b) The correlation length associated with C_s^z in the Haldane phase ($j > j_{c5}$) as a function of j for $\Delta = 1.5$. We plot the data setting $j_{c5} = 0.183$. Fitting the data to the form, $\xi \sim |j - j_{c5}|^\nu$, we estimate the exponent ν .

Fig. 9. The r -dependence of the spin correlation function $C_s^z(r)$ for $j = 1.0$ on a semi-log plot. For $\Delta \leq 1.6$ (the DH phase), $C_s^z(r)$ is divided by the oscillating factor $f(r) = \cos(Qr)$, while it is divided by $f(r) = \exp(i\pi[r/2])$ for $\Delta \geq 2.0$ (the DN phase). The number of kept states is $m = 180$ (for $\Delta = 2.0, 2.5$), and $m = 260$ (for $\Delta = 1.2, 1.4, 1.6$). The numerical errors of the data are smaller than the symbols. Arrows in the figures represent the extrapolated $r = \infty$ values.




Fig.1 T. Hikihara et al.

(a) spin correlation




fig.2(a) T. Hikihara et al.

(b) chiral correlation




Fig.2(b) T. Hikihara et al.

(c) string correlation




Fig.2(c) T. Hikihara et al.




Fig.3 T. Hikihara et al.




Fig.4 T. Hikiyara et al.




Fig.5 T. Hikihara et al.




Fig.6 T. Hikihara et al.




Fig.7 T. Hikihara et al.




Fig.8(a) T. Hikihara et al.




Fig.8(b) T. Hikihara et al.




Fig.9 T. Hikihara et al.